

10/559153

REC'D 16 JUL 2004

WIPO PCT

P1 1193432

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

July 13, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM  
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK  
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT  
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A  
FILING DATE.

APPLICATION NUMBER: 60/475,938

FILING DATE: June 04, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/17175

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS



*M. Tarver*

M. TARVER  
Certifying Officer

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

06-06-03

60475938 OFFICIAL  
SUBSTITUTE for Provisional Application for Patent Cover Sheet PTO/SB/16 (10-01)  
Approved for use through 10/31/2002. OMB 0651-0032  
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

## PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

17611 U.S. PRO  
06/04/0316447 U.S. PRO  
60/475938  
06/04/03

DOCKET NUMBER						21414PV	
INVENTOR(S)							
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)			
Nigel J. Christopher F. David A. John A.		Liverton Claiborne Claremon McCauley		Harleysville, PA 19438 Cambridge, MA 02138 Maple Glen, PA 19002 Maple Glen, PA 19002			
<input type="checkbox"/> Additional inventors are being named on the separately numbered sheets attached hereto							
TITLE OF THE INVENTION (500 characters max)							
N-SUBSTITUTED NONARYL-HETEROCYCLIC NMDA/NR2B ANTAGONISTS							
CORRESPONDENCE ADDRESS							
Direct all Correspondence to: Merck & Co., Inc. Patent Department - RY60-30 P.O. Box 2000 Rahway							
STATE		New Jersey		ZIP CODE		07065	
COUNTRY		U.S.A.					
ENCLOSED APPLICATION PARTS (check all that apply)							
<input checked="" type="checkbox"/> Specification		Number of Pages		44		<input type="checkbox"/> CD(s), Number	
<input type="checkbox"/> Drawing(s)		Number of Sheets				<input type="checkbox"/> Other (specify)	
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76							
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)							
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees						FILING FEE AMOUNT (\$)	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 13-2755						\$160.00	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE

Date 06/04/2003

TYPED or PRINTED NAME Mitul I. Desai

REGISTRATION NO.

46,661

TELEPHONE 732-594-3190

(if appropriate)

EXPRESS MAIL CERTIFICATE	
DATE OF DEPOSIT	June 4, 2003
EXPRESS MAIL NO.	EL 920633267 US
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231.	
MAILED BY	Don Schepisi
DATE	June 4, 2003

In Duplicate

Computer generated form "Transmittal Form-PV" (Application Filing Folder) Merck &amp; Co., Inc., 10/05/2001

TITLE OF THE INVENTION

N-SUBSTITUTED NONARYL-HETEROCYCLIC NMDA/NR2B ANTAGONISTS

FIELD OF THE INVENTION

- 5                    This invention relates to *N*-substituted nonarylheterocyclic compounds. In particular, this invention relates to *N*-substituted nonarylheterocyclic compounds that are NMDA NR2B antagonists useful for the treatment of Parkinson's disease and pain.

10    BACKGROUND OF THE INVENTION


- Ions such as glutamate play a key role in processes related to chronic pain and pain-associated neurotoxicity – primarily by acting through N-methyl-D-aspartate (“NMDA”) receptors. Thus, inhibition of such action – by employing ion channel antagonists, particularly NMDA antagonists – can be beneficial in the treatment and control of Parkinson's disease and pain.

- 15                    NMDA receptors are heteromeric assemblies of subunits, of which two major subunit families designated NR1 and NR2 have been cloned. Without being bound by theory, it is generally believed that the various functional NMDA receptors in the mammalian central nervous system (“CNS”) are only formed by combinations of NR1 and NR2 subunits, which respectively express glycine and glutamate recognition sites. The NR2 subunit family is in turn divided into four individual subunit types: NR2A, NR2B, NR2C, and NR2D. T. Ishii, et al., *J. Biol. Chem.*, 268:2836-2843 (1993), and D.J. Laurie et al., *Mol. Brain Res.*, 51:23-32 (1997) describe how the various resulting combinations produce a variety of NMDA receptors differing in physiological and pharmacological properties such as ion gating properties, magnesium sensitivity, pharmacological profile, as well as in anatomical distribution.

- 20                    For example, while NR1 is found throughout the brain, NR2 subunits are differentially distributed. In particular, it is believed that the distribution map for NR2B lowers the probability of side effects while treating Parkinson's disease or pain. Thus, it would be desirable to provide novel NMDA antagonists that target the NR2B receptor.

SUMMARY OF THE INVENTION

The present invention relates to *N*-substituted nonarylheterocyclic compounds represented by Formula (I):



5

(I)


or pharmaceutically acceptable salts thereof. The present invention also provides pharmaceutical compositions comprising the instant compounds. This invention further provides methods to treat and prevent conditions, including Parkinson's disease, pain, Alzheimer's disease and epilepsy, utilizing the present compounds and compositions.

10

DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are represented by Formula (I):

15



(I)

or pharmaceutically acceptable salts thereof, wherein

HetAr is a 5 or 6 membered heteroaromatic ring containing 1 or 2  
nitrogen ring atoms, or thiazolyl, or thiadiazolyl;  
R<sup>1</sup> and R<sup>2</sup> are independently H, C<sub>1-4</sub>alkyl, fluoro, chloro, bromo, or  
iodo;

A is a bond or -C<sub>1-2</sub>alkyl-; and

20

B is aryl(CH<sub>2</sub>)<sub>0-3</sub>-O-C(O)-, indanyl(CH<sub>2</sub>)<sub>0-3</sub>-O-C(O)-, aryl(CH<sub>2</sub>)<sub>1-3</sub>-C(O)-, aryl-cyclopropyl-C(O)-, aryl(CH<sub>2</sub>)<sub>1-3</sub>-NH-C(O)-, wherein any of the aryl is optionally substituted by 1-5 substituents, each substituent independently is C<sub>1-4</sub>alkyl, fluoro, or chloro.

5

In one aspect, the compounds of this invention are represented by Formula (I), or pharmaceutically acceptable salts thereof, wherein

HetAr is a 6 membered heteroaromatic ring containing 1 or 2 nitrogen ring atoms.

10

In an embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom.

15

In another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms.

20

In a second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is thiazolyl or thiadiazolyl.

25

In an embodiment of this second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is 1,2,4-thiadiazolyl.

30

In another embodiment of this second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is thiazolyl.

In a third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein A is a bond or  $-C_{1-2}alkyl-$ .

5

In an embodiment of this third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein A is a bond.

10

In an other embodiment of this third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

A is methylene.

15

In a further embodiment of this third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

A is  $-C_2 alkyl-$ .

20

In a fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

B is aryl-cyclopropyl- $C(O)-$  or aryl $(CH_2)_{0-3}-O-C(O)-$ , wherein the aryl is optionally substituted with  $C_{1-4}alkyl$ .

25

In an embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

B is aryl-cyclopropyl- $C(O)-$ , wherein the aryl is phenyl, optionally substituted with  $C_{1-4}alkyl$ .

30

In another embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

B is aryl $(CH_2)_{0-3}-O-C(O)-$ , wherein the aryl is phenyl, optionally substituted with  $C_{1-4}alkyl$ .

35

In a further embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

5           B is aryl-O-C(O)-, wherein the aryl is phenyl, optionally substituted with C<sub>1-4</sub>alkyl.

In a still further embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

10           B is aryl(CH<sub>2</sub>)-O-C(O)-, wherein the aryl is phenyl, optionally substituted with C<sub>1-4</sub>alkyl.

In yet another embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

15           B is aryl(CH<sub>2</sub>)-O-C(O)-, wherein the aryl is 4-tolyl.

In a fifth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

20           HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;

A is methylene; and

B is aryl(CH<sub>2</sub>)-O-C(O)-, wherein the aryl is 4-tolyl.

25

In a sixth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

HetAr is 1,2,4-thiadiazolyl;

A is methylene; and

30           B is aryl(CH<sub>2</sub>)-O-C(O)-, wherein the aryl is 4-tolyl.

In a seventh aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

35           HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;

21414PV

A is methylene;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is phenyl, optionally substituted with C<sub>1-4</sub>alkyl.

5           As used herein, "alkyl" as well as other terms having the prefix "alk" such as, for example, alkoxy, alkanoyl, alkenyl, alkynyl and the like, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl and the like.

10           The term "aryl", unless specifically stated otherwise, includes optionally substituted multiple and single ring systems such as, for example, phenyl, naphthyl and tolyl.

          The term "HetAr" includes, for example, heteroaromatic rings such as pyrimidine and pyridine.

15           The term "(CH<sub>2</sub>)<sub>0</sub>" means that the methyl is not present. Thus, "(CH<sub>2</sub>)<sub>0-3</sub>" means that there are from none to three methyls present – that is, three, two, one, or no methyl present. When no methyl groups are present in a linking alkyl group, the link is a direct bond.

          The term "optionally substituted" is intended to include both  
20   substituted and unsubstituted. Thus, for example, optionally substituted aryl can represent a pentafluorophenyl or a phenyl ring. Further, the substitution can be made at any of the groups. For example, substituted aryl(C<sub>1-6</sub>)alkyl includes substitution on the aryl group as well as substitution on the alkyl group.

          Compounds described herein may contain one or more asymmetric  
25   centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and pharmaceutically acceptable salts thereof. The above Formula I is shown without a definitive stereochemistry at certain positions. The present invention includes all  
30   stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Further, mixtures of stereoisomers as well as isolated specific stereoisomers are also included. During the course of the synthetic procedures used to prepare such compounds, or in using racemization or epimerization procedures known to those skilled in the art, the products of such procedures can be a mixture of stereoisomers.



The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (and/or pharmaceutically acceptable salt(s) thereof) as an active ingredient, a pharmaceutically acceptable carrier, and, optionally, other therapeutic ingredients or adjuvants. The instant compositions include those suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

In practice, the compounds represented by Formula I, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient. Further, the compositions can be presented as a powder, as granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid, as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compound represented by Formula I, and/or pharmaceutically acceptable salt(s) thereof, may also be administered by controlled release means and/or delivery devices. The compositions may be prepared by any of the methods of pharmacy. In general, such methods include a step of bringing into association the active ingredient with the carrier that constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and

intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

Thus, the pharmaceutical compositions of this invention may include a  
5 pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable salt of Formula I. The compounds of Formula I, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

The pharmaceutical carrier employed can be, for example, a solid,  
10 liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

In preparing the compositions for oral dosage form, any convenient  
15 pharmaceutical media may be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like may be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like may be used to form oral solid preparations  
20 such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques

A tablet containing the composition of this invention may be prepared  
25 by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of  
30 the powdered compound moistened with an inert liquid diluent. Each tablet preferably contains from about 0.5mg to about 5g of the active ingredient and each cachet or capsule preferably containing from about 0.5mg to about 5g of the active ingredient.

The pharmaceutical compositions of the present invention comprise a  
35 compound represented by Formula I (or pharmaceutically acceptable salts thereof) as

an active ingredient, a pharmaceutically acceptable carrier, and optionally one or more additional therapeutic agents or adjuvants. The instant compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

Pharmaceutical compositions of the present invention suitable for parenteral administration may be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

Pharmaceutical compositions of the present invention suitable for injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability. The pharmaceutical compositions must be stable under the conditions of manufacture and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

Pharmaceutical compositions of the present invention can be in a form suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, or the like. Further, the compositions can be in a form suitable for use in transdermal devices. These formulations may be prepared, utilizing a compound represented by Formula I of this invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the

21414PV

mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

- 5 In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above may include, as appropriate, one or more additional carrier ingredients such as diluents, buffers, flavoring agents, binders, surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation  
10 isotonic with the blood of the intended recipient. Compositions containing a compound described by Formula I, and/or pharmaceutically acceptable salts thereof, may also be prepared in powder or liquid concentrate form.

## ASSAYS

15

### Cell-Based functional assay to determine the $IC_{50}$ of NR2B antagonists

- The ability of selected compounds to inhibit NR1a/NR2B NMDA receptor, as measured by NR1a/NR2B receptor-mediated  $Ca^{2+}$  influx, was assessed by  
20 the following calcium flux assay procedure:

- NR1a/NR2B receptor transfected L(tk-) cells were plated in 96-well format at  $3 \times 10^4$  cells per well and grown for one to two days in normal growth medium (Dulbeccos MEM with Na pyruvate, 4500 mg glucose, pen/strep, glutamine, 10% FCS and 0.5 mg/mL geneticin). NR1a/NR2B-expression in these cells was  
25 induced by the addition of 4-20 nM dexamethasone in the presence of 500  $\mu$ M ketamine for 16 - 24 hours. Solutions of NR2B antagonists were prepared in DMSO and serially diluted with DMSO to yield 10 solutions differing by 3-fold in concentration. A 96-well drug plate was prepared by diluting the DMSO solution 250-fold into assay buffer (Hanks Balanced Salt Solution (HBSS)  $Mg^{2+}$  free (Gibco #14175-079) containing 20 mM HEPES, 2 mM  $CaCl_2$ , 0.1 % BSA and 250  $\mu$ M  
30 Probenecid (Sigma # P-8761)). After induction, the cells were washed twice (Labsystem cell washer, 3 fold dilutions leaving 100  $\mu$ L) with assay buffer and loaded with 4  $\mu$ M of the calcium fluorescence indicator fluo-3 AM (Molecular Probes # P-1241) in assay buffer containing Pluronic F-127 (Molecular Probes # P-3000) and 10  
35  $\mu$ M ketamine at 37  $^{\circ}C$  for one hour. The cells were then washed eight times with

assay buffer leaving 100  $\mu$ L of buffer in each well. Fluorescence intensity was immediately measured in a FLIPR (Fluorometric Imaging Plate Reader, Molecular Devices) using an excitation of 488 nm and emission at 530 nm. Five seconds after starting the recording of fluorescence intensity, 50  $\mu$ L of agonist solution (40  $\mu$ M glutamate /glycine, the final concentration 10  $\mu$ M) was added and after one minute, when fluorescence signal was stable, 50  $\mu$ L of NR2B antagonists and control solutions from the drug plate were added and the fluorescence intensity recorded for another 30 minutes. The  $IC_{50}$  values were determined by a non-linear least squares fitting of the endpoint fluorescence values to Equation #1 below.

10

Equation #1:

$$\text{Endpoint Florescence} = \frac{(Y_{\max} - Y_{\min})}{1 + ([\text{Drug}] / IC_{50})^{nH}} + Y_{\min}$$

15

where,  $Y_{\min}$  is average endpoint fluorescence of the control wells containing 1  $\mu$ M of AMD-2 and  $Y_{\max}$  is the average endpoint fluorescence of wells containing 0.1% DMSO in assay buffer.

20

#### **Binding assay to determine the $K_i$ NR2B Antagonists**

The radioligand binding assay was performed at room temperature in 96-well microtiter plates with a final assay volume of 1.0 mL in 20 mM Hepes buffer (pH 7.4) containing 150 mM NaCl. Solutions of NR2B antagonists were prepared in DMSO and serially diluted with DMSO to yield 20  $\mu$ L of each of 10 solutions differing by 3-fold in concentration. Non-specific binding (NSB) was assessed using AMD-1 (10  $\mu$ M final concentration), and total binding (TB) was measured by addition of DMSO (2% final concentration). Membranes expressing NR1a/NR2B receptors (40 pM final concentration) and tritiated AMD-2 (1 nM final concentration) were added to all wells of the microtiter plate. After 3 hours of incubation at room temperature, samples are filtered through Packard GF/B filters (presoaked in 0.05% PEL, polyethylenimine Sigma P-3143) and washed 10 times with 1 mL of cold 20 mM Hepes buffer per wash. After vacuum drying of the filter plates, 40  $\mu$ L of Packard Microscint-20 was added and bound radioactivity determined in a Packard TopCount. The apparent

35

dissociation constant ( $K_i$ ), the maximum percentage inhibition ( $\%I_{max}$ ), the minimum percentage inhibition ( $\%I_{min}$ ) and the hill slope ( $nH$ ) were determined by a non-linear least squares fitting the bound radioactivity (CPM bound) to Equation #2 below.

5

Equation #2:


$$\text{CPM Bound} = \frac{(\text{SB})(\%I_{max} - \%I_{min})/100}{(1 + ([\text{Drug}]/(K_i (1 + [\text{AMD-2}]/K_D)))^{nH})} + \text{NSB} + (\text{SB})(100 - \%I_{max})/100$$

10

15


where,  $K_D$  is the apparent dissociation constant for the radioligand for the receptor as determined by a hot saturation experiment and SB is the specifically bound radioactivity determined from the difference of TB and NSB control wells.

AMD-1



20

AMD-2




25

AMD-1 can be synthesized according to the general procedure described by C. F. Claiborne et al (Bioorganic & Medchem Letters 13, 697-700 (2003))

The precursor **3** for the synthesis of radiolabelled AMD-1 can be synthesized in accordance with the following procedure:


SCHEME 1a



- 5 In accordance with Scheme 1a, hydrogen chloride is bubbled through a solution of cinnamionitrile **1** in methanol at room temperature. The volatiles are removed under reduced pressure and the resulting residue is triturated with ether and filtered to yield the intermediate imidate **2**. Imidate **2** is dissolved in methanol at ambient temperature, treated with amine **4** (commercially available from Acros Chemicals) at ambient temperature and stirred under argon. The volatiles are removed under reduced pressure and the residue purified by preparative HPLC or
- 10 trituration with ether to afford amidine **3**.

15 Synthesis of Tritiated AMD-2

Scheme 2




- 20 Tritiated AMD-2 was prepared by the following procedure, illustrated in Scheme 2: The precursor **3** (2mg, 0.008mmol) dissolved in dimethylformamide (0.6mL) and potassium carbonate (1.2mg) for 1h. High specific activity tritiated methyl iodide (50mCi, 0.0006mmol, in toluene 1mL, commercially available from American Radiolabeled Chemicals) was added at room temperature and stirred for 2
- 25 hours. The reaction mixture was filtered using a Whatman PTFE 0.45µm syringeless

filter device to remove any insoluble potassium carbonate, washed with Abs. ethanol (2mL, commercially available from Pharmco), and the combined filtrates were concentrated to dryness at room temperature using a rotary evaporator; this also removed any unreacted tritiated methyl iodide. The residue was purified by HPLC chromatography on a Phenomenx Luna C8 semi-prep column (Luna 5 micro C8(2), 250x10.0 mm) using a gradient system of 20/80 acetonitrile/water with 0.1% trifluoroacetic acid to 100% acetonitrile with 0.1% trifluoroacetic acid in 20min. Total activity of the product was 8mCi. Further purification was effected by absorption onto a Waters C-18 Sep-pak column (Waters Sep-Pak PLUS C18) and elution with water followed by absolute ethanol. The product was diluted with absolute ethanol (10mL) before submission for final analysis.

#### Synthesis of unlabelled AMD-2

SCHEME 1b



In accordance with Scheme 1b, hydrogen chloride is bubbled through a solution of cinnamionitrile 1 in methanol at room temperature. The volatiles are removed under reduced pressure and the resulting residue is triturated with ether and filtered to yield the intermediate imidate 2. Imidate 2 is dissolved in methanol at ambient temperature, treated with amine 6 at ambient temperature and stirred under argon. The volatiles are removed under reduced pressure and the residue purified by preparative HPLC or trituration with ether to afford amidine 5.

The compounds of this invention exhibit  $IC_{50}$  and  $K_i$  values of less than 50  $\mu M$  in the functional and binding assays, respectively. It is advantageous that the  $IC_{50}$  and  $K_i$  values be less than 5  $\mu M$  in the functional and binding assays, respectively. It is more advantageous that the  $IC_{50}$  and  $K_i$  values be less than 1  $\mu M$  in the functional and binding assays, respectively. It is still more advantageous that the  $IC_{50}$  and  $K_i$  values be less than 0.1  $\mu M$  in the functional and binding assays, respectively.



The present compounds are NMDA NR2B receptor antagonists, and as such are useful for the treatment and prophylaxis of diseases and disorders mediated through the NR2B receptor. Such diseases and disorders include, but are not limited to, Parkinson's disease, neuropathic pain (such as postherpetic neuralgia, nerve injury, the "dynias", e.g., vulvodynia, phantom limb pain, root avulsions, painful diabetic neuropathy, painful traumatic mononeuropathy, painful polyneuropathy), central pain syndromes (potentially caused by virtually any lesion at any level of the nervous system), and postsurgical pain syndromes (eg, postmastectomy syndrome, postthoracotomy syndrome, stump pain)), bone and joint pain (osteoarthritis), repetitive motion pain, dental pain, cancer pain, myofascial pain (muscular injury, fibromyalgia), perioperative pain (general surgery, gynecological), chronic pain, dysmenorrhea, as well as pain associated with angina, and inflammatory pain of varied origins (e.g. osteoarthritis, rheumatoid arthritis, rheumatic disease, tenosynovitis and gout), headache, migraine and cluster headache, depression, anxiety, schizophrenia, stroke, traumatic brain injury, Alzheimer's disease, cerebral ischemia, amyotrophic lateral sclerosis, Huntington's disease, sensorineural hearing loss, tinnitus, glaucoma, neurological damage caused by epileptic seizures or by neurotoxin poisoning or by impairment of glucose and/or oxygen to the brain, vision loss caused by neurodegeneration of the visual pathway, Restless Leg Syndrome, multi-system atrophy, non-vascular headache, primary hyperalgesia, secondary hyperalgesia, primary allodynia, secondary allodynia, or other pain caused by central sensitization. Compounds of formula I may be used to prevent dyskinesias, particularly the side effects accompanying normal doses of L- Dopa. Furthermore, compounds of formula I may be used to decrease tolerance and/or dependence to opioid treatment of pain, and for treatment of withdrawal syndrome of e.g., alcohol, opioids, and cocaine.

It is understood that compounds of this invention can be administered at prophylactically effective dosage levels to prevent the above-recited conditions, as well as to prevent other conditions mediated through the NMDA NR2B receptor.

30

Compounds of Formula I may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of Formula I. When a

35

compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of Formula I is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of Formula I. Examples of other active ingredients that may be combined with a compound of Formula I, either administered separately or in the same pharmaceutical compositions, include, but are not limited to: (1) non-steroidal anti-inflammatory agents; (2) COX-2 inhibitors; (3) bradykinin B1 receptor antagonists; (4) sodium channel blockers and antagonists; (5) nitric oxide synthase (NOS) inhibitors; (6) glycine site antagonists; (7) potassium channel openers; (8) AMPA/kainate receptor antagonists; (9) calcium channel antagonists; (10) GABA-A receptor modulators (e.g., a GABA-A receptor agonist); (11) matrix metalloprotease (MMP) inhibitors; (12) thrombolytic agents; (13) opioids such as morphine; (14) neutrophil inhibitory factor (NIF); (15) L-Dopa; (16) carbidopa; (17) levodopa/carbidopa; (18) dopamine agonists such as bromocriptine, pergolide, pramipexole, ropinirole; (19) anticholinergics; (20) amantadine; (21) carbidopa; (22) catechol O-methyltransferase ("COMT") inhibitors such as entacapone and tolcapone; (23) Monoamine oxidase B ("MAO-B") inhibitors; (24) opiate agonists or antagonists; (25) 5HT receptor agonists or antagonists; (26) NMDA receptor agonists or antagonists; (27) NK1 antagonists; (28) selective serotonin reuptake inhibitors ("SSRI") and/or selective serotonin and norepinephrine reuptake inhibitors ("SSNRI"); (29) tricyclic antidepressant drugs, (30) norepinephrine modulators; (31) lithium; (32) valproate; and (33) neurontin (gabapentin).

25 Creams, ointments, jellies, solutions, or suspensions containing the instant compounds can be employed for topical use. Mouth washes and gargles are included within the scope of topical use for the purposes of this invention.

30 A formulation intended for the oral administration to humans may conveniently contain from about 0.5mg to about 5g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms can generally contain between from about 1mg to about 1000mg of the active ingredient.

35 The conditions recited herein can be treated or prevented by the administration of from about 0.01mg to about 140mg of the instant compounds per kilogram of body weight per day.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors. Such factors include the age, body weight, general health, sex, and diet of the patient. Other factors include the time and route of administration, rate of excretion, drug combination, and the type and severity of the particular disease undergoing therapy. For example, inflammatory pain may be effectively treated by the administration of from about 0.01mg to about 75mg of the present compound per kilogram of body weight per day, or alternatively about 0.5mg to about 3.5g per patient per day. Neuropathic pain may be effectively treated by the administration of from about 0.01mg to about 125mg of the present compound per kilogram of body weight per day, or alternatively about 0.5mg to about 5.5g per patient per day.

The abbreviations used herein are as follows unless specified otherwise:

15


4-MeBnOH	4-Methylbenzyl alcohol
CDI	1,1'-Carbonyldiimidazole
TEA	Triethylamine
TBSCl	t-Butyldimethylsilyl chloride
DMF	Dimethylformamide
(+)-BINAP	(+)-2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl
NaOtBu	Sodium t-butoxide
DIPEA	Diisopropylethylamine
EtOAc	Ethyl acetate
TBSOTf	t-Butyldimethylsilyl triflate
TBS	t-butyldimethylsilyl
THF	Tetrahydrofuran
DMAP	4-Dimethylaminopyridine
RT	Room temperature
h	Hours
min	Minutes
DCM	Dichloromethane
MeCN	Acetonitrile
iPrOH	2-Propanol
n-BuOH	1-Butanol

EDC 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride  
HOAt 1-Hydroxy-7-azabenzotriazole

## METHODS OF SYNTHESIS


5

## SCHEME 3




10

## SCHEME 4



## SCHEME 5




5

Compounds of the present invention can be prepared according to the Schemes provided below as well as the procedures provided in the Examples. The following Schemes and Examples further describe, but do not limit, the scope of the invention.

10

EXAMPLES 1 AND 2


## EXAMPLE 1



15

(3S,4R)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate

## EXAMPLE 2




- 5 (3R,4S)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-  
carboxylate

## Step 1

## Preparation of 4-Methylbenzyl 4-oxopiperidine-1-carboxylate:

10



- 4-Methylbenzyl alcohol (37.6 g, 308 mmol) was added to a solution of  
1,1'-carbonyldiimidazole (50.0 g, 308 mmol) in DMF at RT and stirred for 1 h. 4-  
15 Piperidone hydrate hydrochloride (commercially available from Sigma-Aldrich, 47.0  
g, 308 mmol) was added, resulting in a reaction mixture that was then heated to 50 °C  
and stirred for 15 h. The reaction mixture was diluted with EtOAc and washed with  
0.1 M HCl, H<sub>2</sub>O (four times), and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated.  
Purification by silica gel chromatography (step gradient elution: 10%, 25%, 50%  
20 EtOAc in hexanes) produced the title compound (42.4 g, 85% yield) as a clear oil.


<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 (d, 2 H), 7.15 (d, 2 H), 5.08 (s, 2 H), 3.79 (t, 4 H),  
2.45 (br s, 4 H) 2.31 (s, 3 H) ppm;

HRMS (ES) *m/z* 248.1281 [(M+H)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>18</sub>NO<sub>3</sub>: 248.1287];

- 25 Anal. C<sub>14</sub>H<sub>17</sub>NO<sub>3</sub>: C, 68.03; H, 7.05; N, 5.59. Found: C, 68.00; H, 6.93; N, 5.66.

## Step 2

## Preparation of (±)-4-methylbenzyl 3-fluoro-4-oxopiperidine-1-carboxylate:



5

A solution of 4-methylbenzyl 4-oxopiperidine-1-carboxylate (21.2 g, 85.7 mmol) and diisopropylethylamine (71.3 mL, 428 mmol) in dichloromethane (425 mL) was cooled to 0 °C and stirred. TBSOTf (29.5 mL, 129 mmol) was added slowly, maintaining the internal temperature below 5 °C. Aqueous NaHCO<sub>3</sub> (20 mL) was added and the layers were separated. The organic layer was washed with NaHCO<sub>3</sub>, H<sub>2</sub>O (two times), and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the crude TBS enol ether.

10

The crude TBS enol ether was dissolved in DMF (125 mL) at RT. Selectfluor® reagent (commercially available from Air Products and Chemicals, Inc., 30.4 g, 85.7 mmol) was added and the reaction mixture was stirred for 10 min. The reaction mixture was partitioned between EtOAc and H<sub>2</sub>O and the organic layer was washed with H<sub>2</sub>O (three times). The combined aqueous layers were extracted with EtOAc (two times) and the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The entire reaction above was repeated and the resulting reaction products were combined to give the title compound (40 g), which was used in the next step without purification. NMR and mass spectral data suggest the ketone functionality in the product exists as a hydrate.

15

20


<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 (m, 2 H), 7.19 (m, 2 H), 5.18 (s, 2 H), 4.81 (br d, 1 H), 4.50 (br d, 1 H), 4.23 (d, 1 H), 3.90 (m, 1 H), 3.60 (m, 1 H), 3.35 (t, 1 H), 2.58 (m, 2 H), 2.35 (s, 3 H) ppm;

25

HRMS (ES) *m/z* 284.1292 [(M+H)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>18</sub>FNO<sub>4</sub>; 284.1293];

Anal. C<sub>14</sub>H<sub>18</sub>FNO<sub>4</sub>•1.2 H<sub>2</sub>O: C, 58.61; H, 6.46; N, 4.88. Found: C, 58.28; H, 6.06; N, 4.72.

30


**Step 3****Preparation of:**

5

(±)-4-methylbenzyl (E)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-carboxylate

and

10



(±)-4-methylbenzyl (Z)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-carboxylate

15

To a solution of (±)-4-methylbenzyl 3-fluoro-4-oxopiperidine-1-carboxylate (40 g, 150 mmol) in toluene (200 mL) at RT was added (carbethoxymethylene)triphenylphosphorane (63.0 g, 181 mmol) and the reaction mixture stirred for 1 h. The reaction mixture was concentrated and purified by silica gel chromatography (gradient elution: 10% to 20% EtOAc in hexanes) to give the olefins (±)-4-methylbenzyl (E)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-carboxylate and (±)-4-methylbenzyl (Z)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-carboxylate (41.0 g, 78% yield, 3 steps) as a 3:1 E:Z mixture. This mixture was utilized directly in the next step. A small sample of the mixture was separated by silica gel chromatography for characterization purposes.



(±)-4-methylbenzyl (E)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-carboxylate: white solid,

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.26 (d, 2 H), 7.17 (d, 2 H), 5.98 (s, 1 H), 5.11 (s, 2 H), 4.85 (m, 1 H), 4.18 (q, 2 H), 4.08 (br d, 1 H), 3.70 (m, 1 H), 3.55 (m, 1 H), 3.41 (m, 1 H), 3.33 (m, 1 H), 2.63 (br d, 1 H), 2.35 (s, 3 H), 1.29 (t, 3 H) ppm;

HRMS (ES) *m/z* 358.1420 [(M+Na)<sup>+</sup>; calcd for C<sub>18</sub>H<sub>22</sub>FNO<sub>4</sub>Na: 358.1425];

Anal. C<sub>18</sub>H<sub>22</sub>FNO<sub>4</sub>: C, 64.21; H, 6.58; N, 4.27. Found: C, 64.46; H, 6.61; N, 4.18.

(±)-4-methylbenzyl (Z)-4-(2-ethoxy-2-oxoethylidene)-3-fluoropiperidine-1-


carboxylate: white solid,

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 (d, 2 H), 7.15 (d, 2 H), 6.41 (m, 1 H), 5.82 (s, 1 H), 5.11 (d, 2 H), 4.61 (m, 1 H), 4.38 (br d, 1 H), 4.16 (q, 2 H), 3.05-2.95 (m, 1 H), 2.9-2.75 (m, 2 H), 2.33 (s, 3 H), 2.13 (m, 1 H), 1.27 (t, 3 H) ppm;

HRMS (ES) *m/z* 358.1422 [(M+Na)<sup>+</sup>; calcd for C<sub>18</sub>H<sub>22</sub>FNO<sub>4</sub>Na: 358.1425].


#### Step 4:

Preparation of:



(±)-cis 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate

and



(±)-trans 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate

To a solution of the olefin mixture from **Step 3** (10.0 g, 29.8 mmol) in toluene (160 mL) and  $\text{CH}_2\text{Cl}_2$  (120 mL) was added diphenylsilane (5.53 mL, 29.8 mmol) and (R)-BINAP (1.86 g, 2.98 mmol). Sodium t-butoxide (0.29 g, 2.98 mmol) and CuCl (0.30 g, 2.98 mmol) were then added, the reaction mixture was protected from light and stirred for 15 h. Additional portions of diphenylsilane (2.76 mL), NaOtBu (0.29 g) and CuCl (0.30 g) were added and the reaction mixture was stirred at RT for 24h. The mixture was then filtered through celite and concentrated. Purification on silica gel (step gradient elution: 5%, 10%, 15%, 25%, 30% EtOAc in hexanes) gave recovered starting materials (3.5 g, 35% yield), ( $\pm$ )-cis 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate (5.0 g, 50% yield) and ( $\pm$ )-trans 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate (1.2 g, 12% yield).

15 ( $\pm$ )-cis 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate: clear oil,

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.25 (d, 2 H), 7.15 (d, 2 H), 5.10 (s, 2 H), 4.80-4.20 (m, 3 H), 4.15 (q, 2 H), 3.10-2.73 (m, 2 H), 2.52 (dd, 1 H), 2.35 (s, 3 H), 2.30 (dd, 1 H), 2.10 (m, 1 H), 1.72-1.48 (m, 2 H), 1.29 (t, 3 H) ppm;

20 HRMS (ES)  $m/z$  338.1689  $[(\text{M}+\text{H})^+]$ ; calcd for  $\text{C}_{18}\text{H}_{25}\text{FNO}_4$ : 338.1762].


( $\pm$ )-trans 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate: clear oil,

25  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24 (d, 2 H), 7.15 (d, 2 H), 5.08 (s, 2 H), 4.50-3.95 (m, 3 H), 4.15 (q, 2 H), 2.81 (br t, 2 H), 2.70 (br d, 1 H), 2.35 (s, 3 H), 2.17 (m, 2 H), 1.89 (br d, 1 H), 1.25 (m, 1 H), 1.22 (t, 3 H) ppm;

HRMS (ES)  $m/z$  338.1699  $[(\text{M}+\text{H})^+]$ ; calcd for  $\text{C}_{18}\text{H}_{25}\text{FNO}_4$ : 338.1762].

**Step 5**

**Preparation of (±)-((cis)-3-fluoro-1-[(4-methylbenzyl)oxy]carbonyl)piperidin-4-yl)acetic acid:**



5


To a solution of (±)-cis 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate (10.0 g, 29.6 mmol) in THF (50 mL) was added aqueous NaOH (1M, 50 mL). The reaction mixture was stirred at RT for 5 h and then  
10 diluted with EtOAc and 1M HCl. The layers were separated and the aqueous extracted with EtOAc twice. The combined organics were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the title compound (9.1 g) as a white solid which was used in the next step without further purification.

15 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 (d, 2 H), 7.15 (d, 2 H), 5.08 (s, 2 H), 4.79-4.16 (m, 3 H), 3.05-2.75 (m, 2 H), 2.59 (dd, 1 H), 2.36 (dd, 1 H), 2.31 (s, 3 H), 2.20-2.02 (m, 1 H), 1.60 (m, 2 H) ppm;  
HRMS (ES) *m/z* 310.1457 [(M+H)<sup>+</sup>; calcd for C<sub>16</sub>H<sub>21</sub>FNO<sub>4</sub>: 310.1449].  
Anal. C<sub>16</sub>H<sub>20</sub>FNO<sub>4</sub>•0.15 H<sub>2</sub>O: C, 62.13; H, 6.52; N, 4.53. Found: C, 61.55; H, 6.37; N,  
20 4.41.

**Step 6**

**Preparation of (±)-cis-4-methylbenzyl 4-(aminomethyl)-3-fluoropiperidine-1-carboxylate:**

25




- To a suspension of crude acid ( $\pm$ )-((cis)-3-fluoro-1-{[(4-methylbenzyl)oxy]carbonyl}piperidin-4-yl)acetic acid (9.1 g, 29.4 mmol) in toluene (80 mL) was added triethylamine (10.2 mL, 73.5 mmol) and diphenylphosphoryl azide (9.52 mL, 44.1 mmol). The reaction mixture was heated to 70 °C and stirred for 20 min. A mixture of dioxane (80 mL) and 1 M NaOH (80 mL) was added and the reaction mixture was cooled to RT. The reaction mixture was concentrated to remove the dioxane and extracted with EtOAc three times, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The residue was suspended in  $\text{CH}_2\text{Cl}_2$ , stirred for 30 min, and the white precipitate filtered off. The filtrate was concentrated to give crude product (7.5 g) as a yellow oil, used directly in the next step. An analytical sample was purified by silica gel chromatography (gradient elution:  $\text{CH}_2\text{Cl}_2$  to 80:20:2  $\text{CH}_2\text{Cl}_2$  : MeOH :  $\text{NH}_4\text{OH}$ ) for characterization:

- $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24 (d, 2 H), 7.15 (d, 2 H), 5.08 (s, 2 H), 4.90-4.18 (m, 3 H), 2.95-2.75 (m, 2 H), 2.79 (dd, 1 H), 2.70 (dd, 1 H), 2.35 (s, 3 H), 1.59 (m, 3 H) ppm;  
HRMS (ES)  $m/z$  281.1658 [ $(\text{M}+\text{H})^+$ ; calcd for  $\text{C}_{15}\text{H}_{22}\text{FN}_2\text{O}_2$ : 281.1660].


#### Step 7

- 20 **Preparation of:**



- 25 (3S,4R)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate

and



(3R,4S)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate

5 Two sealed tubes were each charged with a mixture of crude ( $\pm$ )-cis-4-methylbenzyl 4-(aminomethyl)-3-fluoropiperidine-1-carboxylate (Step 6, 3.7 g, 13.2 mmol) and 2-chloropyrimidine (1.51 g, 13.2 mmol) in n-butanol/diisopropyl-ethylamine (1:1, 13 mL). The tubes were sealed and the mixtures heated to 140 °C and stirred for 2 h. After cooling to RT, the reaction mixtures were combined and  
10 diluted with EtOAc and sat NaHCO<sub>3</sub>. The layers were separated and the organic was washed with H<sub>2</sub>O and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification by silica gel chromatography (gradient elution: 1:1 hexanes:EtOAc to EtOAc) gave racemic cis-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate (6.9 g, 65% yield, 3 steps) as a white solid.  
15

The enantiomers were separated by preparative HPLC on a ChiralPak AD column (5 cm x 50 cm, 20 $\mu$ M) with MeOH:MeCN (15:85, 150 mL/min) as eluant. The HCl salt of **Example 1** was prepared by dissolving (3S,4R)-cis-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate (6.9  
20 g, 19.3 mmol) in iPrOH (100 mL) at 65 °C. A solution of HCl in iPrOH (1.608 M, 12.6 mL, 20.2 mmol) was added and the solution was slowly cooled to RT over 15 h. Et<sub>2</sub>O (25 mL) was added, the mixture stirred for 3h, cooled to 0 °C, stirred for 1h and filtered to give (3S,4R)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate hydrochloride as a white solid (7.0 g, 92%  
25 recovery).

The hydrochloride salt of (3R,4S)-4-methylbenzyl-3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate was prepared using a similar  
30 procedure.

**(3S,4R)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate•HCl:**

$[\alpha]_D -36.4^\circ$  (c 0.17, MeOH);

5 Melting Point 149-150 °C;

$^1\text{H NMR}$  (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.58 (br s, 2 H), 7.21 (d, 2 H), 7.17 (d, 2 H), 6.99 (t, 1 H), 5.06 (s, 2 H), 4.79 (m, 1 H), 4.42 (t, 1 H), 4.21 (d, 1 H), 3.60 (dd, 1 H), 3.50 (dd, 1 H), 3.15-2.80 (m, 2 H), 2.30 (s, 3 H), 2.10 (m, 1 H), 1.61 (m, 2 H) ppm;  
HRMS (ES)  $m/z$  359.1879  $[(M+H)^+]$ ; calcd for  $\text{C}_{19}\text{H}_{24}\text{FN}_4\text{O}_2$ : 359.1878];

10 Anal.  $\text{C}_{19}\text{H}_{23}\text{FN}_4\text{O}_2 \cdot \text{HCl} \cdot 0.2 \text{H}_2\text{O}$ : C, 57.27; H, 6.17; N, 14.06. Found: C, 57.22; H, 6.37; N, 14.16.

**(3R,4S)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate •HCl:**

15  $[\alpha]_D +34.9^\circ$  (c 0.18, MeOH);

Melting Point 149-150 °C;


$^1\text{H NMR}$  (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.58 (br s, 2 H), 7.21 (d, 2 H), 7.17 (d, 2 H), 6.99 (t, 1 H), 5.06 (s, 2 H), 4.79 (m, 1 H), 4.42 (t, 1 H), 4.21 (d, 1 H), 3.60 (dd, 1 H), 3.50 (dd, 1 H), 3.15-2.80 (m, 2 H), 2.30 (s, 3 H), 2.10 (m, 1 H), 1.61 (m, 2 H) ppm;

20 HRMS (ES)  $m/z$  359.1870  $[(M+H)^+]$ ; calcd for  $\text{C}_{19}\text{H}_{24}\text{FN}_4\text{O}_2$ : 359.1878].

Anal.  $\text{C}_{19}\text{H}_{23}\text{FN}_4\text{O}_2 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$ : C, 56.50; H, 6.24; N, 13.87. Found: C, 56.68; H, 6.27; N, 13.80.

**EXAMPLES 3 and 4**

25



**(-)-trans-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate (Example 3) and (+)-trans-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-**

30 **ylamino)methyl]piperidine-1-carboxylate (Example 4)**

The title compounds were prepared from (±)-trans 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate (Examples 1 and 2, Step 4), utilizing the procedures described in Steps 5, 6 and 7 from Examples 1 and 2.

5 (-)-trans-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate •HCl:

$[\alpha]_D -11.5^\circ$  (c 0.22, MeOH);

Melting Point 113-114°C;

10  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.80-8.39 (m, 2 H), 7.24 (d, 2 H), 7.15 (d, 2 H), 7.00 (t, 1 H), 5.08 (s, 2 H), 4.49-4.21 (m, 2 H), 4.00 (d, 1 H), 3.81 (dd, 1H), 3.58 (dd, 1 H), 2.95 (m, 1 H), 2.31 (s, 3 H), 2.12 (m, 1 H), 1.90 (m, 1 H), 1.34 (m, 1 H) ppm;

HRMS (ES)  $m/z$  359.1867  $[(M+H)^+]$ ; calcd for  $\text{C}_{19}\text{H}_{24}\text{FN}_4\text{O}_2$ : 359.1878].

Anal.  $\text{C}_{19}\text{H}_{23}\text{FN}_4\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$ : C, 55.27; H, 6.35; N, 13.57. Found: C, 55.08; H, 6.11; N, 13.36.

15

(+)-trans-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate •HCl:

$[\alpha]_D +16.3^\circ$  (c 0.17, MeOH);

Melting Point 113-114°C

20  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.80-8.39 (m, 2 H), 7.24 (d, 2 H), 7.15 (d, 2 H), 7.00 (t, 1 H), 5.08 (s, 2 H), 4.49-4.21 (m, 2 H), 4.00 (d, 1 H), 3.81 (dd, 1H), 3.58 (dd, 1 H), 2.95 (m, 1 H), 2.31 (s, 3 H), 2.12 (m, 1 H), 1.90 (m, 1 H), 1.34 (m, 1 H) ppm;


HRMS (ES)  $m/z$  359.1873  $[(M+H)^+]$ ; calcd for  $\text{C}_{19}\text{H}_{24}\text{FN}_4\text{O}_2$ : 359.1878];

Anal.  $\text{C}_{19}\text{H}_{23}\text{FN}_4\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$ : C, 55.27; H, 6.35; N, 13.57. Found: C, 55.18; H, 6.11;

25 N, 13.38.


## EXAMPLE 5

Preparation of:



5

(3S,4R)-N-[(3-cis-fluoro-1-[(1R,2R)-2-phenylcyclopropyl]carbonyl]piperidin-4-yl)methylpyrimidin-2-amine



10

(3S,4R)-4-methylbenzyl 3-fluoro-4-[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate (Example 1) (400 mg, 1.16 mmol) was dissolved in 3N HCl (4 mL), heated to 90 °C and stirred for 1h. The reaction mixture was concentrated and the crude amine hydrochloride dissolved in DMF (3 mL) at RT and (1R,2R)-2-phenylcyclopropanecarboxylic acid (188 mg, 1.16 mmol) (prepared as described by T. N. Riley and C. G. Brier, *J. Org. Chem*, 15, 1187-1188, 1972), EDC (222 mg, 1.16 mmol), HOAt (158 mg, 1.16 mmol) and triethylamine (323 µL, 2.32 mmol) were added. The reaction mixture was heated to 70 °C and stirred for 10 min. After cooling to RT, the reaction mixture was diluted with EtOAc and H<sub>2</sub>O. The layers were separated and the organic was washed with H<sub>2</sub>O and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification by silica gel chromatography (gradient elution: 1:1 hexanes:EtOAc to 90:10:1 EtOAc:MeOH:NH<sub>4</sub>OH) gave the title compound (320 mg, 78% yield) as a white solid. The compound was characterized as the hydrochloride salt, which was prepared in a similar manner to that described in Example 1, Step 7.



$[\alpha]_D -164^\circ$  (c 0.22, MeOH);


Melting Point 127-128°C;

$^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.59 (br s, 2 H), 7.25 (m, 2 H), 7.17 (m, 3 H), 7.01 (t, 1 H), 4.85 (m, 2 H), 4.65-4.30 (m, 2 H), 3.55 (m, 2 H), 3.30 (m, 1 H), 2.80 (m, 1 H), 2.39 (m, 1 H), 2.30-2.10 (m, 2 H), 1.75 (m, 1 H), 1.55 (m, 2 H), 1.33 (m, 1 H) ppm;

HRMS (ES)  $m/z$  355.1925  $[(M+H)^+]$ ; calcd for  $\text{C}_{20}\text{H}_{24}\text{FN}_4\text{O}$ : 355.1929];

Anal.  $\text{C}_{20}\text{H}_{24}\text{FN}_4\text{O} \cdot \text{HCl} \cdot \text{H}_2\text{O}$ : C, 58.75; H, 6.41; N, 13.70. Found: C, 58.57; H, 6.44; N, 13.54.


# EXAMPLE 6



15 (-)-cis-4-methylbenzyl 3-fluoro-4-[(1,3,4-thiadiazol-2-ylamino)methyl]piperidine-1-carboxylate

## Step 1

Preparation of:



20 (±)-cis-4-methylbenzyl 4-[(hydrazinocarbonothioyl)amino]methyl-3-fluoropiperidine-1-carboxylate

25 A solution of (±)-cis-4-methylbenzyl 4-(aminomethyl)-3-fluoropiperidine-1-carboxylate (Example 1, Step 6) (200 mg, 0.71 mmol) in DMF (2

10

## 15 Step 2

20

30

**Melting Point 115-117°C;**


3

- 32 -

HRMS (ES)  $m/z$  365.1437 [(M+H)<sup>+</sup>; calcd for C<sub>17</sub>H<sub>22</sub>FN<sub>4</sub>O<sub>2</sub>S: 365.1442].  
Anal. C<sub>17</sub>H<sub>21</sub>FN<sub>4</sub>O<sub>2</sub>S: C, 56.03; H, 5.81; N, 15.37. Found: C, 55.82; H, 5.74; N, 15.11.

### EXAMPLE 7

5



(+)-trans-4-methylbenzyl 3-fluoro-4-[(1,3,4-thiadiazol-2-ylamino)methyl]piperidine-1-carboxylate:

10

The title compound was prepared from (±)-trans 4-methylbenzyl 4-(2-ethoxy-2-oxoethyl)-3-fluoropiperidine-1-carboxylate (Examples 1 and 2, Step 4), utilizing the procedures described in Examples 1 and 2, Steps 5 and 6, followed by thiadiazole formation as described in Example 6. The (+)-enantiomer was separated by chiral HPLC as described in Example 1 and 2 Step 7.

15

[α]<sub>D</sub> +37.1° (c 0.56, MeOH);

Melting Point 93-95 °C;

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.36 (s, 1 H), 7.28 (d, 2 H), 7.19 (d, 2 H), 6.56 (br s, 1 H), 5.04 (s, 2 H), 4.45 (m, 2 H), 4.17 (m, 1 H), 3.56 (m, 2 H), 2.80 (m, 2 H), 2.35 (s, 3 H), 2.10 (m, 1 H), 1.96 (m, 1 H), 1.39 (m, 1 H) ppm;

20

HRMS (ES)  $m/z$  365.1432 [(M+H)<sup>+</sup>; calcd for C<sub>17</sub>H<sub>22</sub>FN<sub>4</sub>O<sub>2</sub>S: 365.1442];


Anal. C<sub>17</sub>H<sub>21</sub>FN<sub>4</sub>O<sub>2</sub>S•0.25 H<sub>2</sub>O: C, 55.34; H, 5.87; N, 15.19. Found: C, 55.50; H, 5.61; N, 14.81.

25

21414PV

## WHAT IS CLAIMED IS:

1. A compound having the formula (I):



5

(I)

or a pharmaceutically acceptable salt thereof, wherein

HetAr is a 5 or 6 membered heteroaromatic ring containing 1 or 2 nitrogen ring atoms, thiazolyl, or thiadiazolyl;

- 10 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C<sub>1-4</sub>alkyl, fluoro, chloro, bromo, or iodo;

A is a bond or -C<sub>1-2</sub>alkyl-; and

- 15 B is aryl(CH<sub>2</sub>)<sub>0-3</sub>-O-C(O)-, indanyl(CH<sub>2</sub>)<sub>0-3</sub>-O-C(O)-, aryl(CH<sub>2</sub>)<sub>1-3</sub>-C(O)-, aryl-cyclopropyl-C(O)-, aryl(CH<sub>2</sub>)<sub>1-3</sub>-NH-C(O)-, wherein any of the aryl is optionally substituted by 1-5 substituents, each substituent independently is C<sub>1-4</sub>alkyl, fluoro, or chloro.

2. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

- 20 HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom.

3. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

- 25 HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms.

4. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
HetAr is thiazolyl.
- 5 5. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
HetAr is thiadiazolyl.
- 10 6. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
HetAr is 1,2,4 thiadiazolyl.
- 15 7. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
A is a bond.
- 20 8. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
A is methylene.
- 25 9. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
A is -C<sub>2</sub> alkyl-.
10. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
B is aryl-cyclopropyl-C(O)-, wherein said aryl is optionally substituted as defined in Claim 1.
- 30 11. The compound according to Claim 10, or a pharmaceutically acceptable salt thereof, wherein said aryl is phenyl, optionally substituted as defined in Claim 1.

21414PV


12. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
B is  $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$ , wherein said aryl is optionally substituted as defined in Claim 1.
- 5
13. The compound according to Claim 1, or a pharmaceutically acceptable salts thereof, wherein  
B is  $\text{aryl}(\text{CH}_2)-\text{O}-\text{C}(\text{O})-$ , wherein said aryl is optionally substituted as defined in Claim 1.
- 10
14. The compound according to Claim 13, or a pharmaceutically acceptable salt thereof, wherein said aryl is optionally substituted with  $\text{C}_{1-4}$  alkyl.
- 15
15. The compound according to Claim 14, or a pharmaceutically acceptable salt thereof, wherein said aryl is 4-tolyl.
16. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring  
atoms;  
A is methylene; and  
B is  $\text{aryl}(\text{CH}_2)-\text{O}-\text{C}(\text{O})-$ , wherein said aryl is optionally substituted as defined in Claim 1.
- 20
17. The compound according to Claim 16, or a pharmaceutically acceptable salt thereof, wherein  
said aryl is optionally substituted with  $\text{C}_{1-4}$  alkyl.
- 25
18. The compound according to Claim 17, or a pharmaceutically acceptable salt thereof, wherein  
said aryl is 4-tolyl.
- 30

21414PV

19. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;  
5 A is methylene; and  
B is 4-tolyl(CH<sub>2</sub>)-O-C(O)-.
20. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
10 HetAr is thiadiazolyl;  
A is methylene; and  
B is aryl(CH<sub>2</sub>)-O-C(O)-, wherein said aryl is optionally substituted as defined in Claim 1.
21. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein  
15 HetAr is 1,2,4-thiadiazolyl;  
A is methylene; and  
B is 4-tolyl(CH<sub>2</sub>)-O-C(O)-.
22. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein  
20 HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;  
A is methylene; and  
25 B is aryl-cyclopropyl-C(O)-.
23. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein  
30 HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms; and  
B is phenyl-cyclopropyl-C(O)-.


21414PV

24. The compound according to Claim 1, wherein said compound is



5 or a pharmaceutically acceptable salt thereof.

25. The compound according to Claim 1, wherein said compound is




10

or a pharmaceutically acceptable salt thereof.



21414PV

26. The compound according to Claim 1, wherein said compound  
is




5

or a pharmaceutically acceptable salt thereof.

27. The compound according to Claim 1, wherein said compound  
is

10




(-)

or a pharmaceutically acceptable salt thereof.

28. The compound according to Claim 1, wherein said compound  
is

15




| k(+)

20 or a pharmaceutically acceptable salt thereof.


21414PV

29. The compound according to Claim 1, wherein said compound  
is




5  
or a pharmaceutically acceptable salt thereof.

30. The compound according to Claim 1, wherein said compound  
is  
10



or a pharmaceutically acceptable salt thereof.

31. The compound according to Claim 1, wherein said compound  
is  
15



20 or a pharmaceutically acceptable salt thereof.

21414PV

32. A pharmaceutical composition comprising an inert carrier and a therapeutically effective amount of a compound according to Claim 1.

33. The pharmaceutical composition according to Claim 32, further comprising a second therapeutic agent selected from the group consisting of: (i) non-steroidal anti-inflammatory agents; (ii) COX-2 inhibitors; (iii) bradykinin B1 receptor antagonists; (iv) sodium channel blockers and antagonists; (v) nitric oxide synthase (NOS) inhibitors; (vi) glycine site antagonists; (vii) potassium channel openers; (viii) AMPA/kainate receptor antagonists; (ix) calcium channel antagonists; (x) GABA-A receptor modulators (e.g., a GABA-A receptor agonist); (xi) matrix metalloprotease (MMP) inhibitors; (xii) thrombolytic agents; (xiii) opioids such as morphine; (xiv) neutrophil inhibitory factor (NIF); (xv) L-Dopa; (xvi) carbidopa; (xvii) levodopa/carbidopa; (xviii) dopamine agonists such as bromocriptine, pergolide, pramipexole, ropinirole; (xix) anticholinergics; (xx) amantadine; (xxi) carbidopa; (xxii) catechol O-methyltransferase ("COMT") inhibitors such as entacapone and tolcapone; (xxiii) Monoamine oxidase B ("MAO-B") inhibitors; (xiv) opiate agonists or antagonists; (xv) 5HT receptor agonists or antagonists; (xvi) NMDA receptor agonists or antagonists; (xvii) NK1 antagonists; (xviii) selective serotonin reuptake inhibitors ("SSRI") and/or selective serotonin and norepinephrine reuptake inhibitors ("SSNRI"); (xxix) tricyclic antidepressant drugs, (xxx) norepinephrine modulators; (xxxii) lithium; (xxxiii) valproate; and (xxxiii) neurontin (gabapentin).

34. The pharmaceutical composition according to Claim 32 useful for the treatment of pain, migraine, schizophrenia, depression, anxiety, cluster headache, Alzheimer's disease, stroke, or epilepsy.

35. The pharmaceutical composition according to Claim 33 useful for the treatment of pain, migraine, schizophrenia, depression, anxiety, cluster headache, Alzheimer's disease, stroke, or epilepsy.

36. The pharmaceutical composition according to claim 32 useful for the treatment of Parkinson's disease.

37. The pharmaceutical composition according to claim 33 useful for the treatment of Parkinson's disease.

38. A method for treating or preventing pain in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

39. A method for treating or preventing migraine, cluster headache, depression, anxiety, schizophrenia, Alzheimer's disease, or stroke in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

40. A method for treating or preventing Parkinson's disease in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

41. A method for treating or preventing chronic, visceral, inflammatory and neuropathic pain syndromes in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

42. A method for treating or preventing pain resulting from, or associated with, traumatic nerve injury, nerve compression or entrapment, postherpetic neuralgia, trigeminal neuralgia, diabetic neuropathy, cancer and chemotherapy, in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

43. A method for treating or preventing chronic lower back pain in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

21414PV

44. A method for treating or preventing phantom limb pain in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

5

45. A method for treating or preventing HIV- and HIV treatment-induced neuropathy, chronic pelvic pain, neuroma pain, complex regional pain syndrome, chronic arthritic pain and related neuralgias in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

10

46. A method for treating or preventing epilepsy and partial and generalized tonic seizures in a patient in need thereof comprising administering to said patient a therapeutically effective amount, or a prophylactically effective amount, of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

15


TITLE OF THE INVENTION

N-SUBSTITUTED NONARYL-HETEROCYCLIC NMDA/NR2B ANTAGONISTS

ABSTRACT OF THE DISCLOSURE

5

Compounds represented by Formula (I):



(I)

- 10 or pharmaceutically acceptable salts thereof, are effective as NMDA NR2B antagonists useful for treating conditions such as, for example, Parkinson's disease, Alzheimer's disease, migraine, epilepsy and pain.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**